



1. Find, with proof, all integers  $n$  such that  $2^6 + 2^9 + 2^n$  is the square of an integer.
2. Find, with proof, all real numbers  $a$  such that  $|x - 1| - |x - 2| + |x - 4| = a$  has exactly 3 solutions.
3. As Lisa hurried to copy down the last problem of her math homework assignment at the end of class, she got as far as

$$0 = 9x^8 - 28x^6 -$$



## SOLUTIONS

1. If  $m^2 = 2^6 + 2^9 + 2^n = 576 + 2^n = 24^2 + 2^n$ , then  $2^n = m^2 - 24^2 = (m - 24)(m + 24)$ . Therefore,  $m + 24$  and  $m - 24$  must each be powers of 2. Let  $m + 24 = 2^k$  and  $m - 24 = 2^p$  where,  $p < k$  and  $p + k = n$ . Then  $2^k - 2^p = 48$  which implies  $2^p$  divides 48, so that  $p \leq 4$ . Trying  $p = 0, 1, 2, 3, 4$  gives  $p = 4, k = 6$  and  $n = 10$  as the only possible value for  $n$ .

2. We begin by graphing the function  $y = f(x) = |x - 1| - |x - 2| + |x - 4|$ . If  $x \leq 1$ , then  $x - 1 \leq 0, x - 2 \leq 0$  and  $x - 4 \leq 0$ , so we have

$$y = -(x - 1) + (x - 2) - (x - 4) = -x + 3.$$

If  $1 \leq x \leq 2$ , then  $x - 1 \geq 0, x - 2 \leq 0$  and  $x - 4 \leq 0$ , so

$$y = (x - 1) + (x - 2) - (x - 4) = x + 1.$$

In the interval  $2 \leq x \leq 4$  we have  $x - 1 \geq 0, x - 2 \geq 0$  but  $x - 4 \leq 0$ , so

$$y = (x - 1) - (x - 2) - (x - 4) = -x + 5.$$

Finally, for  $x \geq 4$  we have

$$y = (x - 1) - (x - 2) + (x - 4) = x - 3.$$

By piecing together the relevant parts of these four linear functions, we get the graph of the function  $f(x)$  shown at the right.

Thus, for the original equation to have exactly three solutions, we have to choose  $a$  so that the horizontal line  $y = a$  touches the graph of  $f(x)$  at exactly three points. From the graph, this happens only for  $a = 2$  and  $a = 3$ .

3.  $P(x) = 9x^8 - 28x^6 - \dots$  Since the coefficient of  $x^7$  is zero, the sum of the roots of  $P(x) = 0$  is zero. Let the seven identical roots be  $a$  and the desired eighth root be  $b$ . Then  $7a + b = 0$  or  $b = -7a$ . Standardizing  $P(x)$ , the sum of the products of the roots taken two at a time is  $-\frac{28}{9}$ . Therefore,  $C^2 = 7 \cdot \frac{28}{9}$

4. a. Let the desired pair-square set be  $\{2012, a, b\}$ . Since  $2012 + 13 = 2025 = 45^2$ , try  $a = 13$  as a second member of the set. Then

$$2012 + b = x^2 \quad \text{and} \quad 13 + b = y^2 \quad \text{for some integers } x \text{ and } y.$$

Subtracting these two equations gives  $1999 = x^2 - y^2 = (x + y)(x - y)$ . Since 1999 is a prime number,  $x + y = 1999$  and  $x - y = 1$ . From these two equations we obtain  $x = 1000$ . Then  $b = 1000^2 - 2012 = 1000000 - 2012 = 997988$ . Thus, one desired pair-square set of size 3 is  $\{13, 2012, 997988\}$ .

(Note:  $\{292, 2012, 45077\}$  and  $\{488, 2012, 143912\}$  also work. There are others.)

- b. Every integer has one of the four forms  $4k$ ;  $4k + 1$ ;  $4k + 2$  and  $4k + 3$  for integers  $k$ .

First we prove that the square of an integer must have one of the forms  $4n$  or  $4n + 1$ .

Proof:

$$(i) \quad (4k)^2 = 4(4k^2) = 4n$$

$$(ii) \quad (4k + 1)^2 = 16k^2 + 8k + 1 = 4(4k^2 + 2k) + 1 = 4n + 1$$

$$(iii) \quad (4k + 2)^2 = 16k^2 + 16k + 4 = 4(4k^2 + 4k + 1) = 4n$$

$$(iv) \quad (4k + 3)^2 = 16k^2 + 24k + 9 = 4(4k^2 + 6k + 2) + 1 = 4n + 1$$

Therefore, the square of an integer must have one of the forms  $4n$  or  $4n + 1$ .

Let  $S$  be a pair-square set of size 3. Suppose that the set  $S$  contains the two odd numbers  $a$  and  $b$ . Since  $a + b$  is an even square, it must have form  $4n$ , and therefore  $a$  and  $b$  cannot both have form  $4k + 1$ , nor can they both have form  $4k + 3$ . It follows that we can write  $a = 4k + 1$  and  $b = 4k + 3$ .

We derive a contradiction by showing that there is no possibility for the third member  $z$  of  $S$ . Indeed, if  $z$  has form  $4k$  or  $4k + 3$ , then  $z + b$  is not a square, and if  $z$  has form  $4k + 2$  then  $z + a$  is not a square.

5. It is easy to show that  $\triangle ADP$  is isosceles (note the marked congruent angles).

Thus  $DP = 2012$  and all we need is the length of  $PC$ .

Since  $\triangle ADP$  and  $\triangle CBQ$  are congruent isosceles triangles,

$\angle PAB \cong \angle DAP \cong \angle CQB$ , making  $\overline{AP}$  parallel to  $\overline{QC}$ .

Similarly, the other two angle bisectors are parallel, making  $EFGH$  a parallelogram.

Since  $\angle DAB$  and  $\angle ADC$  are supplementary,  $m\angle DAB + m\angle ADC = 180$ .

Since  $m$